

Effects of Inoculants (*Chlorobium limicola* and *Rhodopseudomonas palustris*) on Nutrient Uptake and Growth in Cucumber

Shuang Shao^{1,*}, Hongyao Zhao¹, Xiaolei Guo², Lijie Guan¹ and Haiyan Li¹

¹College of Pharmaceutical and Biological Engineering, Shenyang University of Chemical Technology, 110142, Liaoning, China

²Liaoning Provincial Seed Administration, 110034, Liaoning, China

Abstract: Rhizobacteria is a prosperous for promoting plant growth for the superiority of reducing environmental damages. Two Strains of *Chlorobium limicola* and *Rhodopseudomonas palustris* were supplied in the experiment as potential inoculants for cucumber. Significant enhancement of the availability of macronutrient elements N, P and K were observed in soil, and further improvement on the uptake of them was also obtained in cucumber plants. Accumulation of essential micronutrients of Fe and Zn were detected both in roots and in shoots. The two stains increased chlorophyll and carotenoid synthesis, plant height, stem diameter, wet weight and dry weight. Various dose has significantly effect on plant growth stimulation, *C. Limicola* with 10^7 cells mL⁻¹ and *R. Palustris* with 10^8 cells mL⁻¹ seem to be better on the whole.

Keywords: Biofertilizer, cucumber, growth, nutrien.

INTRODUCTION

Utilization of fertilizers is really necessary for plant growth, but excessive and repeated use of chemical fertilizers may spoil the soil, water and even atmosphere [1]. Rhizobacteria containing biofertilizers are perfect tools to improve plant nutrient and production and reduce environmental damages [2]. Nitrogen, phosphorus and potassium are major limiting factors to plant growth. Plant growth promoting rhizobacteria (PGPR) including N- fixe, phosphate and potassium solubilizing bacteria (PSB and KSB) stimulate the growth of plants by helping to provide nutrient to the host plant, or indirectly by positively influencing root growth and morphology or by aiding other beneficial symbiotic relationships, or helping to control pathogenic organism [3, 4]. *Chlorobium limicola* and *Rhodopseudomonas palustris* are the diverse group of photosynthetic prokaryotes growing frequently in soils, which are known to fix atmospheric nitrogen[5] and induce resistance, seem to achieve their promised contribution to plant growth. Micronutrient elements are essential to both plant growth and human health. About one third people in the world suffer from the inadequate intakes of Fe, Zn and vitamin A [6], especially in developing countries [7]. Cucumber (*Cucumis sativus* L.) cultivation has an important role in agricultural production, especially under protection. Nutrients are required for maximum growth rate of cucumber. And the optimal weight proportions among the nutrients

maybe is that, in relation to N = 100, close to 75 K, 13 P, 9 Ca, and 9 Mg. In present study, we checked the role of the two strains in macronutrient and micronutrient elements uptake and growth promotion of cucumber.

MATERIALS AND METHODS

Chlorobium limicola and *Rhodopseudomonas palustris* are two newly isolated strains of photosynthetic bacteria. Seeds of cucumber (*Cucumis sativus* L. cv. *Jinchun4*) were surface sterilized with HgCl₂ (0.1%) for 10 min, rinsed with deionized water and placed on filter paper moistured by deionized water, and germinated at 28°C in constant temperature incubator in dark. After germination the seedlings were transferred to plastic pot ($\Phi_{\text{bottom}}10\text{cm}\times16\text{cm}\times\Phi_{\text{top}}15\text{cm}$) containing 1.2 kg autoclaved soil. The cells in cul-tured bacterial broth of both two strains were adjusted to 10^7 , 10^8 and 10^9 cells mL⁻¹, and 150 mL of bacterial broth was applied to each pot every two days, and the application of deionized water and chemical fertilizer were using as controls. The experiment of examining cucumber growth was set in a Completely Randomized Block Design with eight treatments each content four replicates and the replicate represented by ten plants under controlled conditions at $26 \pm 2^\circ\text{C}$ and a 16 hr photoperiod created by using illumination incubator. The morphological characters such as height of the plants, stem diameter, wet weight and dry weight were esti-mated at 40 days after treatment are calculated. We used the Minolta SPAD-502 chlorophyll meter, ± 1.0 SPAD (arbitrary units), to measure the relative chlorophyll contents and chromatography for carotenoid contents of the third fully developed leaves.

*Address correspondence to this author at the College of Pharmaceutical and Biological Engineering, Shenyang University of Chemical Technology, 110142, Liaoning, China; Tel: 86-024-13840034285; Fax: 86-024-89385582; E-mail: shaoshuang403@163.com

Shoot and root tissues were separated after harvesting and were air-dried at 70°C for 5 days, and soil samples were collected after the experiment and air-dried for chemical analysis. Plants were digested in H₂SO₄-H₂O₂ for the determination of total nitrogen. Available N in soils was determined by alkaline-proliferation law. P was determined by the Berthelot reaction and molybdenum blue method [8], soil samples were extracted with 0.5 M NaHCO₃ (pH 8.5). Plants and soils were digested using 1M NH₄OAC (pH 7) for the determination of K using an atomic absorption spectrometer. Atomic absorption spectrometer was also used for the determination of Fe and Zn [9].

The data was analyzed statistically for standard deviation by using sigma plot software. The mean values were compared, using Duncan's multiple range test at P<0.05.

RESULTS

The treatment inoculation with *C. Limicola* and *R. Palustris* significantly increased N, P and K uptake in

cucumber plants. The two strains had stronger effect on N uptake then P and K uptake. The amounts of available N, P and K nutrients in soils were significantly increased as compared to control, which might be attribute to the capability to fix atmospheric nitrogen, solubilize phosphate and potassium by *C. Limicola* and *R. Palustris*. *R. Palustris* with 10⁸ cells mL⁻¹ and *C. limicola* with 10⁷ cells mL⁻¹ have even stronger and more stable effect then with other concentrations (Table 1).

Accumulation of Fe in both plant parts of roots and shoots increased significantly in various treatments of *C. Limicola* and *R. Palustris* as compared to the control of deionized water, and even higher then the control of chemical fertilizer in roots, and reaching to the maximum level in *R. Palustris* with 10⁸ cells mL⁻¹. Level of Zn showed similar trend with that of Fe except reaching to the maximum level in *C. Limicola* with 10⁷ cells mL⁻¹ in roots and *R. Palustris* with 10⁷ cells mL⁻¹ in shoots. It is obviously that the two stains could promote the enrich of Fe and Zn in cucumber, and the effect seems to be stronger in roots than in shoots (Table 2).

Table 1: Effects of *C. Limicola* and *R. Palustris* on Macronutrient Uptake of Cucumber and on the Available N, P, K Concentrations in the Soils (n=4; ± s.e.)

Treatment (cells mL ⁻¹)	cucumber(mg.g dw ⁻¹)			soils(mg kg ⁻¹)		
	N	P	K	N	P	K
deionized water	19.1±0.82c	1.83±0.10b	27.5±2.41b	70.12±4.32c	8.92±0.50d	80.22±7.8c
<i>R. palustris</i> (10 ⁷)	22.3±0.78ab	2.09±0.08a	31.6±2.01a	92.75±5.19a	10.89±0.67ab	98.07±8.5b
<i>R. palustris</i> (10 ⁸)	22.4±0.98a	2.06±0.11a	31.5±1.93a	98.85±2.07a	10.65±0.81ab	110.16±10.5a
<i>R. palustris</i> (10 ⁹)	20.0±1.11bc	2.04±0.08a	30.5±2.21a	88.80±3.55ab	10.31±0.15bc	98.97±8.7b
<i>C. limicola</i> (10 ⁷)	21.8±0.83ab	2.10±0.10a	31.0±1.58a	82.63±4.25ab	11.33±0.38a	112.34±5.6a
<i>C. limicola</i> (10 ⁸)	20.4±1.02bc	2.12±0.14a	30.2±1.33a	90.76±1.77ab	11.50±1.16a	94.61±7.2b
<i>C. limicola</i> (10 ⁹)	21.6±1.30ab	2.08±0.09a	30.6±2.71a	81.74±1.84b	9.62±1.27cd	95.32±5.8b

Table 2: Effect of *C. Limicola* and *R. Palustris* on the Contents of Fe and Zn in Cucumber (n=4; ± s.e.) (mg. kg dw⁻¹)

Treatment (cells mL ⁻¹)	roots		shoots	
	Fe	Zn	Fe	Zn
deionized water	415.6±18.3e	123.1±3.21d	532.7±19.5c	146.6±4.68c
chemical fertilizer	475.5±17.6d	144.6±4.44c	608.9±20.2abc	176.2±3.11ab
<i>R. palustris</i> (10 ⁷)	601.4±16.4ab	156.8±2.61b	612.8±21.8ab	182.7±4.65a
<i>R. palustris</i> (10 ⁸)	655.4±20.5a	166.2±0.98ab	654.4±16.4a	175.5±4.12ab
<i>R. palustris</i> (10 ⁹)	602.8±17.5ab	167.5±5.21ab	572.7±17.2bc	174.2±2.08ab
<i>C. limicola</i> (10 ⁷)	588.2±21.4bc	174.9±3.89a	594.6±18.5abc	178.4±1.22ab
<i>C. limicola</i> (10 ⁸)	532.3±18.2c	162.8±4.56ab	566.2±18.9bc	175.9±5.97ab
<i>C. limicola</i> (10 ⁹)	560.1±18.6bc	170.8±2.17a	548.3±17.7c	165.8±3.26b

Table 3: Effects of *C. Limicola* and *R. palustris* on the Plant Height, Stem Diameter, Wet Weight, Dry Weight, Relative Chlorophyll Contents and Carotinoid Contents of Cucumber (n=4;±s.e.)

Treatment (cells mL ⁻¹)	Height (cm plant ⁻¹)	stem diameter (cm plant ⁻¹)	wet weight (g plant ⁻¹)	dry weight (g plant ⁻¹)	Chlorophyll (mg g ⁻¹)	Carotinoid (mg g ⁻¹)
deionized water	13.8±0.86e	1.37±0.05d	7.37±0.84e	0.72±0.11c	10.0±0.82d	3.25±0.48b
chemical fertilizer	14.1±1.29e	1.40±0.07cd	7.51±0.62de	0.74±0.07c	10.6±0.98c	2.99±0.71c
<i>R. palustris</i> (10 ⁷)	25.6±0.76a	1.43±0.06bcd	9.98±0.91a	0.88±0.02a	11.1±0.89b	3.55±0.27a
<i>R. palustris</i> (10 ⁸)	25.0±0.84a	1.43±0.16bcd	9.98±0.01a	0.88±0.22a	11.8±0.52a	3.64±0.67a
<i>R. palustris</i> (10 ⁹)	15.0±1.28d	1.47±0.14bc	7.96±0.92cd	0.61±0.01d	6.50±0.99e	1.80±0.20d
<i>C. limicola</i> (10 ⁷)	23.1±0.42b	1.50±0.10ab	8.10±0.62c	0.62±0.02d	11.6±1.38ab	1.89±0.28d
<i>C. limicola</i> (10 ⁸)	18.2±1.36c	1.57±0.03a	9.05±0.61b	0.71±0.08c	11.2±1.35b	3.57±0.71a
<i>C. limicola</i> (10 ⁹)	21.5±0.46b	1.57±0.22a	10.0±0.673a	0.80±0.06b	10.1±0.74d	3.26±0.09b

Both strains showed a promoting effect on plant growth, morphological and biochemical parameters representing at Table 3 showed a best response to the *R. Palustris* inoculants with 10⁸ cells mL⁻¹, and treatment with 10⁷ cells mL⁻¹ is following. Morphological parameters showed a good response to *C. limicola* with 10⁹ cells mL⁻¹, however, several parameters responding to *R. Palustris* with 10⁸ cells mL⁻¹ were lower than control (Table 3).

DISCUSSION

The experiment demonstrated that *C. Limicola* and *R. Palustris* stimulated the availability of the macronutrient element N, P and K in soils, and make further improvement on the uptake of them in cucumber plants. The two strains could also promote the accumulation of essential micronutrients of Fe and Zn. This probably due to the effect of bacteria on creating the availability of Fe and Zn. Fe is the maximum micronutrient for plant growth. Fe and Zn can accelerate chlorophyll synthesis. So we can see the application of the two strains enhanced growth, while the probably physiological basis might be that an increased chlorophyll and carotinoid synthesis makes a stronger photosynthesis, photosynthesis may additionally provides energy for nitrogen fixation, allowing for more efficient plant growth [10], therefore, wet weight and dry weight enhanced, and a more intensive uptake of nutrients might also cause a promoting growth of cucumber [11]. The increasing N uptake in our experiment might be related to the fact that photosynthetic bacteria which *C. Limicola* and *R. Palustris* falls into is a genus which fixes atmospheric nitrogen [12]. Whether the increasing uptake of P and K should due to the role of strains by aiding other beneficial symbiotic relationships or their own role

needs further research. The perfect behaviour of the two strains appears therefore a priority to really envisage a future for the use of them as an inoculant for cucumber.

ACKNOWLEDGEMENTS

This research was financially supported by Shenyang University of Chemical Technology and a grant from Shenyang Science and technology bureau (no. F12-108-3-00). The helpful comments of the editor and anonymous reviewers are gratefully acknowledged.

REFERENCES

- [1] Mytton L. Nitrogen Fixation In Institute of Grassland and Environmental Research Rep. Institute of Grassland and Environ. Res. Aberystwyth, UK 1993; p. 46-5.
- [2] Vessey JK. Plant growth promoting rhizobacteria as biofertilizers. *Plant Soil* 2003; 255: 571-16. <http://dx.doi.org/10.1023/A:1026037216893>
- [3] Whipps JM. Microbial interactions and biocontrol in the rhizosphere. *J Exp Bot* 2001; 52: 487-25. http://dx.doi.org/10.1093/jexbot/52.suppl_1.487
- [4] Zehnder GW, Murphy JF, Sikora EJ, Kloepper JW. Application to rhizobacteria for induced resistance. *Eur J Plant Pathol* 2001; 107: 39-12. <http://dx.doi.org/10.1023/A:1008732400383>
- [5] Paul WL, Gary PR. Nitrogen fixation by photosynthetic bacteria. *Photosynth Res* 2002; 73: 115-4. <http://dx.doi.org/10.1023/A:1020497619288>
- [6] Food and Agriculture Organization of the United Nations (FAO). Preventing micronutrient malnutrition. A guide to food-based approaches. International Life Science Institute: Washington DC 1997.
- [7] Miflin B. Crop biotechnology. Where now? *Plant Physiol* 2000; 123: 17-12. <http://dx.doi.org/10.1104/pp.123.1.17>
- [8] Page AL, Mille RH, Keeney DR. *Methods of Soil Analysis: Part 2. Chemical and Microbiological Properties* 2nd ed. Amer Society of Agronomy 1982; 9: 1159.
- [9] Han YS. *The Experimental Technology of Food Chemistry*. Agricultural University. Beijing Chin (in Chinese) 1996.

- [10] Giraud E, Fleischman D. Nitrogen-fixing symbiosis between photosynthetic bacteria and legumes. *Photosynth Res* 2004; 82: 115-6.
<http://dx.doi.org/10.1007/s11120-004-1768-1>
- [11] Lévai L. Bacteria containing fertilizer: A tool for sustainable agriculture. XV. International Plant Nutrition Colloquium, Peking 2005.
- [12] Han HS, Supanjani, Lee KD. Effect of co-inoculation with phosphate and potassium solubilizing bacteria on mineral uptake and growth of pepper and cucumber. *Plant Soil Environ* 2006; 52: 130-7.

Received on 13-02-2014

Accepted on 16-03-2014

Published on 31-03-2014

DOI: <http://dx.doi.org/10.6000/1929-5634.2014.03.01.5>